

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

3ake
126

REPORT OF PROGRESS IN BARBERRY ERADICATION FOR THE CALENDAR YEAR

ENDED DECEMBER 31, 1926

By

F. E. Kempton and Lynn D. Hutton

Reprinted, with different pagination, from the Cereal Courier 19 (4): 49-77,
February 28, 1927.

(NOT FOR PUBLICATION)

CONTENTS

	Page
Introduction -----	1
Organization and Funds -----	2
Cooperation -----	3
Finances -----	3
Investigations -----	3
Stem-Rust Epidemiology Studies -----	4
Classification of barberry species, varieties, and hybrids -----	5
Inoculation of barberry species, varieties, and hybrids with stem rust -----	5, 6
Publicity and Education Activities -----	6, 7, 8
Surveys -----	9
Original Survey -----	9
Second Survey -----	10
Resurvey -----	10
Areas of escaped barberries -----	10, 11
Eradication -----	11
Summarized Results -----	12
Summary for 1926 -----	12
Summary for nine years -----	12
Tables	
1. Original survey, properties, Jan. 1, 1926, to Dec. 31, 1926 ----	13
2. Original survey, bushes and seedlings, Jan. 1, 1926, to Dec. 31, 1926 --	14
3. Original survey, properties, April 1, 1918, to Dec. 31, 1926 ---	15
4. Original survey, bushes and seedlings, April 1, 1918, to Dec. 31, 1926	16
5. Second survey, properties, Jan. 1, 1926, to Dec. 31, 1926 -----	17
6. Second survey, bushes and seedlings, Jan. 1, 1926, to Dec. 31, 1926 --	18
7. Second survey, properties, Jan. 1, 1922, to Dec. 31, 1926 -----	19
8. Second survey, bushes and seedlings, Jan. 1, 1922, to Dec. 31, 1926 --	20
9. Resurvey, properties, Jan. 1, 1926, to Dec. 31, 1926 -----	21
10. Resurvey, sprouting bushes and seedlings, Jan. 1, 1926, to Dec. 31, 1926	22
11. Resurvey, properties, April 1, 1918, to Dec. 31, 1926 -----	23
12. Resurvey, sprouting bushes and seedlings, Apr. 1, 1918, to Dec. 31, 1926	24
13. Chemical treatment, Jan. 1, 1926, to Dec. 31, 1926 -----	25
14. Chemical treatment, Sept. 1, 1921, to Dec. 31, 1926 -----	26
15. Chemicals used, Jan. 1, 1926, to Dec. 31, 1926 -----	27
16. Chemicals used, Sept. 1, 1921, to Dec. 31, 1926 -----	28
17. Grand summary, original bushes, sprouting bushes, and seedlings April 1, 1918, to Dec. 31, 1926 -----	29

REPORT OF PROGRESS IN BARBERRY ERADICATION FOR THE CALENDAR YEAR ENDED
DECEMBER 31, 1926

By

F. E. Kempton, Assoc. Pathologist in Charge, and Lynn D. Hutton, Assoc.
Pathologist

INTRODUCTION

In the nine years that the barberry-eradication campaign has been in progress more than 14,300,000 common barberries have been eradicated. This is a monumental record of plant-pest eradication.

The removal of these bushes has aided materially in reducing stem-rust losses of small grains. In the eastern States of the barberry-eradication area positive control of local epidemics of stem rust has resulted from the removal of the harmful barberries. A material reduction of the stem-rust losses from destructive epidemics in the spring-wheat States has followed the eradication of common barberries. In many localities in these more western States positive control has been effected.

The publicity activities have acquainted the public with the purpose and progress of the campaign and have resulted in gaining its confidence and cooperation. As a result, the owners and occupants of the properties on which barberries were found have given whole-hearted cooperation in survey and eradication. In many instances, valuable services in labor and chemicals for eradication have been given.

Educational activities in the grade schools, high schools, and colleges are being emphasized. The younger generation is being taught to know black stem rust and its relation to the common barberry. As a result of this educational program, future citizens are being trained to know and eradicate the harmful barberries and to prevent their reintroduction.

The results are only temporary, however, unless the campaign is carried to its satisfactory conclusion. A great many more barberries remain in the 13 States which comprise the eradication area. The preliminary survey has not yet reached 57 of the 920 counties it is necessary to survey. Only 209 counties have been covered by the second more intensive survey. Continued resurveys will be necessary to find and eradicate seedlings and sprouting bushes.

ORGANIZATION AND FUNDS

The barberry-eradication campaign is conducted by the Office of Cereal Crops and Diseases, Bureau of Plant Industry, U. S. Department of Agriculture, in cooperation with 13 north-central, grain-growing States. These are Colorado, Illinois, Indiana, Iowa, Michigan, Minnesota, Montana, Nebraska, North Dakota, Ohio, South Dakota, Wisconsin, and Wyoming.

In each State, a State Leader and a corps of field agents carry out the various field operations. Collaborators, agents, and cooperators in the experiment station and extension divisions of the Agricultural Colleges and in the State Departments of Agriculture, and other interested organizations, aid in the eradication, investigational, and educational activities.

In 1926, there were used 46 full-time employees and 253 part-time agents in conducting the campaign. The field agents are employed during the summer months. Men with two or more years of college training in agriculture or related subjects usually are selected. Training in plant pathology, agronomy, or botany is required. Special attention is given to procuring men who not only are adequately trained but who have a real interest in the work. They also must have a correct ethical attitude toward scientific accuracy and honest public service. Men specially trained in news writing and publicity are valuable aids to the State Leaders in promoting the educational phases of the campaign. A premium is put on experience and ability. The more mature men with longer experience and proved ability in barberry eradication and in teaching agriculture are retained for several summers when possible.

All field men are required to master the necessary details of the scientific facts concerning stem rust and the common barberry. These men also must be able to identify other rusts of grains and to recognize barberries of all ages and sizes, in any environment. Training is given in methods of procedure in survey and eradication and in making contacts with property owners, officials, and business men.

Supervision in the field is under the direction of the State Leader in each State. An assistant leader aids in field supervision in some States. Squad leaders are placed in charge of small groups and are responsible for survey and eradication in the areas assigned to each squad. If an agent trained in newspaper writing and publicity is available, he promotes the publicity and educational phases under the direction of the State Leader, in cooperation with squad leaders.

Cooperation

Cooperation is had with the State Agricultural Colleges of all the above-named States and the State Departments of Agriculture of most of them. The Conference for the Prevention of Grain Rust provides funds and personnel for aiding in publicity and educational activities and establishing contacts with commercial, agricultural, and State organizations. This Conference is an organization composed of representatives of commercial, agricultural, educational, and investigational organizations of the 13 States.

State and county superintendents of public instruction, professors and instructors in universities, colleges, and normal schools, supervisors of teachers, and the teachers themselves in these States cooperate in the movement to teach the story of stem-rust control by barberry eradication to the rising generation.

Commercial organizations and local farm organizations, county agricultural agents and club leaders, State and county crop reporters, State, county, and township weed-control officials and the thousands of property owners whose properties are inspected for barberries usually cooperate whole-heartedly in the campaign.

Finances

The annual Federal appropriation for the fiscal year ending June 30, 1927, is \$375,000. This is the same as for the past fiscal year. Of this amount, \$75,000 became available only on the certification of contributions totaling an additional \$75,000 by interested agencies in the 13 cooperating States. The total aid furnished in cash and services by State and other agencies for the present fiscal year is larger than the amount required to be certified and is estimated at approximately \$83,000. Federal funds totaling \$2,567,715 have been appropriated and allotted during the period from April 1, 1918, to June 30, 1927. State and other interested agencies have allotted approximately \$652,837 in funds and services. This makes a total of \$3,220,602 allotted to this campaign to June 30, 1927. Approximately \$3,068,000 of Federal and State contributions combined has been expended to December 31, 1926.

INVESTIGATIONS

Three investigational phases of the barberry-eradication campaign are carried forward. These are (1) stem-rust epidemiology studies, (2) classification of barberry species, varieties, and hybrids, and (3) the inoculation of barberry species, varieties, and hybrids with stem rust.

, Stem Rust Epidemiology Studies

Under the supervision of Dr. E. C. Stakman, Agent, University Farm, St. Paul, Minn., stem-rust epidemiology studies are being carried forward each year. These stem-rust studies include the investigation of the sources, time of appearance, prevalence, and spread of stem rust, and estimates of the damage done to small grains. These studies help to determine the aid rendered by barberry eradication in reducing stem-rust losses of small grain. Through these studies, the effect of barberries as local sources of stem rust is traced and the general stem-rust conditions in local areas of each State from year to year are determined. Information also is obtained as to stem-rust conditions in groups of States and in the whole United States.

The early appearance of stem rust on barberries, its subsequent spread to near-by grains and grasses, and its later spread in the form of more wide-spread epidemics are correlated as far as possible. Hundreds of instances of the spread of rust from common barberries to grains and grasses have been recorded. Many have been added during the past year. The members of the barberry-eradication force cooperate closely in this work. State collaborators and cooperators of the barberry-eradication States and of the surrounding States and Canada make observations and furnish data. The Office of Mycology and Disease Survey, Bureau of Plant Industry, U. S. Department of Agriculture, through collaborators in each State, collects and furnishes estimates of the losses from stem rust.

Farmers in many localities give valuable information concerning the occurrence of stem rust each year on individual farms. These reports are investigated. In many instances the harmful barberries have been located and eradicated. Through the reports of severe local outbreaks of stem rust in fields of grain, it has been possible to locate many scattered barberries and seedlings that have grown up along ditches, fence rows, and in near-by woodlands that would have been difficult to find by other means.

Invariably when all barberries are eradicated from local areas, stem rust ceases to be a serious menace to the growing of small grains. It is true that, in some of these areas, a late general spread of stem rust occurs in epidemic years, but the severe damage formerly caused by local barberries does not occur.

Classification of Barberry Species, Varieties, and Hybrids

A *Berberis* collection is being assembled at Bell, Maryland, by Mr. B. Y. Morrison, Associate Horticulturist. The Office of Horticulture and the Office of Cereal Crops and Diseases cooperate in the assembling, growing, and studying of this collection. An attempt is made to include the various species, varieties, and hybrids that are grown or are likely to be grown in the United States. These are studied, described, illustrated, and classified. Plants of species of unknown reaction to stem rust are propagated and sent to St. Paul, Minn., where this susceptibility is studied.

The following is a brief summary of Mr. Morrison's report for 1926:

The work at Bell [Maryland] has been furthered by the cooperation of the Office of Foreign Plant Introduction through which office seed has been secured from the Royal Botanic Gardens at Kew, England, and Edinburgh, Scotland, of all species of barberries in those collections which are not represented already in our collection. Through them seed was obtained also from the Vilmorin Arboretum and from M. Pardé, Director of the Forestry School, at Les Barres, France. The Bell collection now contains 85 good species, 23 dubious species, and 11 hybrids of known parentage.

From the one-year potted plants and the nursery 333 plants of 43 species were sent to University Farm, St. Paul, Minn.

Sheets of illustrations are being prepared showing the life history of the different species from germination to fruiting. In this way it is possible to record very conspicuously the range of variations in the different stages of the plant's life.

The work on the material for the monograph progresses, but is much more complete in some sections of the genus than in others.

Inoculation of Barberry Species, Varieties, and Hybrids with Stem Rust

The common barberry (*Berberis vulgaris* L.) and its horticultural varieties are not the only barberries that may carry and spread stem rust. The barberries known to be susceptible to this disease were listed by E. C. Stakman and M. N. Levine in "A Partial Report on the Susceptibility and Resistance of *Berberis* and Related Genera to Stem Rust," *Cereal Courier* 15: 278-287. September 30, 1923.

The investigation of the susceptibility of other barberries to stem rust is now carried forward by Mr. Ralph U. Cotter. In his report of December 31, 1926, 130 species, varieties, and hybrids are listed as on hand at University Farm, St. Paul, Minn. Of some, there is only one plant; of others, there are several. In 1925, seven species or varieties of barberries were first reported as infected. In 1926, 11 others were reported infected for the first time. The infections in 1926 resulted from 206 series of inoculations of various species and varieties of barberries with stem rust. The ones that failed to become infected on the first trial were inoculated several times, using telial material of different varieties of stem rust collected in different parts of the country.

The studies on the nature of the resistance of barberries to stem rust also are being continued.

PUBLICITY AND EDUCATION ACTIVITIES

Publicity activities always have been an important part of the campaign. Materials designed to inform the public how to identify both the common barberry and the stem rust are a part of all publicity. A special attempt is made to play up items of local importance in each community. Unusual findings of barberries and outstanding examples of the spread of stem rust from barberries are given local and State-wide emphasis. This necessitates an intensive use of news items, demonstrations, exhibits, window displays, radio broadcasts, and talks. Lantern slide series dealing with local and State problems supplement the talks. General and State problems and accomplishments also are shown by news articles, motion pictures, bulletins, circulars, posters, colored plates, and circular letters.

Barberry eradication must be permanent. The future generations should be taught that the common barberry has caused severe damage to small grain in the past and that it must not be allowed to exist or gain a foothold again in grain-growing areas. To accomplish this, the story of stem rust and the barberry is being taught to the rising generation through the public schools. Full cooperation has been volunteered by the State superintendents of public instruction in some States, and the county superintendents of schools are aiding in many counties. State universities, colleges, normal schools, and teachers' colleges are becoming enthusiastic about presenting this story to teachers. Teachers in vocational and other high schools, and in city and country grade schools are being furnished outlines and materials and are cooperating to a large extent in teaching their pupils about stem rust and the common barberry.

A Federal representative of barberry eradication, Mr. Noel F. Thompson, is maintained in the office of the Conference for the Prevention of Grain Rust. This office acts as the clearing house for publicity and education activities and cooperates by furnishing funds, invaluable services, and many materials not otherwise obtainable. Mr. Thompson, in his summary of the publicity and education activities of 1926, reports as follows:

A very satisfying improvement was noted in the publicity and education activities carried on in 1926 over those of preceding years. The results also are becoming increasingly evident, both in the attitude of the public toward the campaign and in the actual aid rendered by individuals whose only contact has been through these activities.

While these activities are correlated through the office of The Conference for the Prevention of Grain Rust, the major portion of the work is carried on directly by the State Leaders, and to them is due full credit for the success of this phase of the barberry-eradication campaign.

The publicity activities are being conducted through newspaper articles, circular letters, demonstrations and exhibits, and radio talks. The preparation of bulletins and circulars and speaking at various public meetings are necessary adjuncts.

The newspaper publicity for the most part was confined to news articles, and these, in general, were given to the local papers in the areas where survey is in progress. Over 1,300 such articles were printed in 1926. In addition, over 120 articles of wider news value were syndicated.

The circular letter, if judiciously used, is very effective publicity. In the past, many of the State Leaders have sent these letters to every farmer in the areas where intensive survey was conducted. These letters, usually accompanied by a bulletin, are sent shortly in advance of the arrival of the field men, paving the way for the survey and making the work much easier and more effective. In 1926, for the first time, all the 13 States used this system, sending out over 147,000 letters.

Letters also are being used to supply up-to-date and reliable information to a number of selected mailing lists. These lists include extension workers, legislators, weed inspectors, prominent and influential farmers, bankers, grain dealers, and other business men. In all, about 40 such lists were used in 1926 and each list was used an average of $2\frac{1}{2}$ times. Many of these lists will be enlarged and improved.

The demonstration is recognized as a valuable publicity feature and is used regularly in all 13 States. All the squads of field men are supplied with window-display outfits and these are placed in the towns of the areas being surveyed. Field demonstrations are staged where a well-defined spread of rust from a barberry is found. Street corner demonstrations are used in the evenings in some localities. Fair demonstrations or exhibits also are used regularly. During 1926, there were 71 field demonstrations, 372 window demonstrations, 150 street corner and miscellaneous demonstrations, and 126 fair demonstrations, making a total of 719 demonstrations for the year.

The radio was used to some extent in 1926. In addition to flashes sent out from Washington, the State Leaders supplied speakers or subject-matter for nine talks on the common barberry and stem rust.

Five new bulletins or circulars written in whole or in part by the State Leaders were published in 1926. Such publications are of special value since they supply detailed information in regard to the particular conditions within the State. They supplement the more general bulletins and circulars printed in Washington.

The State Leaders or their representatives spoke at 535 meetings in 1926.

Educational instruction about the common barberry and stem rust in the schools is becoming one of the major phases of our publicity and educational activities. Efforts are being made to have the story of black stem rust incorporated into the regular curriculum of as many schools as possible and preliminary efforts are meeting with considerable success. Teachers of rural schools, high schools, normal schools, and colleges are being supplied with study material, usually consisting of a lesson plan, colored charts, bulletins, and specimens of rusted barberries and grains. In many cases, these materials come to the public school teacher through her county or State superintendent of schools. During the school year 1925-26, such material was supplied to over 33,000 teachers as compared with 10,000 for the previous year. Attention is now being directed to the normal schools and other teacher-training schools so that the teacher may learn the story of the rust before reaching her school.

A summary of the publicity and education phases of the barberry-eradication campaign would not be complete without mention of the very important part played by the Conference for the Prevention of Grain Rust, an organization of business men, scientists, and farmers interested in the welfare of agriculture. This organization has supplied much of the demonstration material, such as window-display sets and special fair demonstration attractions. They also have furnished nearly all of the colored materials used in 1926, including colored plates, charts, and post-cards. Large numbers of mounted barberry specimens and considerable direct financial aid also have been furnished. Equally important has been the advice and encouragement received from this Conference.

SURVEYS

Three surveys are in progress. These are (1) the original or first survey, (2) the second survey, and (3) the resurveys. The first survey is almost completed. Almost 863 of the 920 counties necessary to survey have been covered. The second, or more intensive, survey is in progress and about 200 counties have been covered. Resurvey for eliminating sprouts and seedlings after the first and second survey is somewhat continuous. Portions of some counties covered by the second survey have been checked for overlooked bushes and seedlings developing since the second survey. It is apparent that additional surveys of some areas will be necessary.

Original Survey

The original survey, prior to 1926, for the most part was a rapid one for the purpose of destroying the greatest number of fruiting barberries in the shortest possible time. By the end of 1925, the original survey had been completed in all of the area except portions of Illinois, Michigan, Montana, and Ohio.

During 1926, the original survey was continued along the more intensive lines which are being used for second survey. Some areas, because of lack of settlement, or arid or forested conditions, may be excepted from survey, or only certain portions may be completely inspected. However, any original survey completed in these areas will be done intensively.

Second Survey

In the second survey, an attempt is made to eliminate every barberry bush and seedling from both city and rural properties. The yard and garden, hedges and fence rows, pasture lands, woodlots, woods, thickets, groves, parks, cemeteries, and all places where barberries may grow, are covered by a complete, intensive strip-by-strip survey. Only such portions of a property are excepted from survey as are actually cultivated, but uncultivated areas within cultivated fields also are inspected.

This method is used in covering all urban and rural properties on or near which fruiting barberries have been found. The history of settlement, type of agriculture, comparative number of barberries first planted, their later spread, the stem-rust history, results of preliminary surveys, and various ecological factors help to determine the extent of this survey.

In recently settled areas, brush lands, timbered flood plains, the cut-over timberlands, mountainous and arid districts, remote woodlands such as National Forests, and the grazing lands may be left for later survey, or not covered at all, as developments show necessary.

Resurvey

Resurvey is the revisiting of the properties from which barberries have been eradicated, in order to eliminate sprouts and to find and destroy seedlings. This activity includes the inspection not only of the property itself but of surrounding properties to which seeds may have been spread. If fruiting bushes have been found on the property or if the information about the planting is meager, the entire property and the surrounding properties are covered intensively for a sufficient distance to insure finding all scattered seedlings. The interval between successive resurveys is such that sprouts, overlooked small bushes, and seedlings will not have had time to produce seeds. This interval usually is three years or longer, depending upon conditions.

Areas of Escaped Barberries

Every property surrounding a fruiting barberry bush is potentially a part of an area of escaped bushes. A complete survey is made of the properties on which fruiting bushes have been found, and of the surrounding properties, to determine whether or not there is a spread of barberries. If either scattered bushes or seedlings are found, the full extent of the spread is determined by further survey.

Full records with maps are made of each area of escaped barberry bushes. In making the survey and mapping each area, definite landmarks are indicated. These are designated on the map with symbols and the same symbols usually are inscribed on the landmarks such as trees, fence posts, and large rocks. This method locates the permanent boundaries of the area as surveyed. If succeeding inspections show further spread of seedlings the added portions are included with additional markings.

One of the principal difficulties in complete eradication of all barberries is the finding and destroying of the scattered seedlings. Fruiting barberry bushes produce thousands of seeds. These seeds are scattered by birds and other agencies to surrounding properties. Just how far they may be carried has not been ascertained. It is certain that they have been carried at least three miles, and possibly further. Some of the seeds grow immediately. Some lie dormant for one to several years before germinating. Some seeds are known to have remained dormant at least seven years.

Seedlings have been found scattered in orchards, groves, woodlots, woodlands, brushy pastures, thickets, swamp lands, river-bottom undergrowth, brushy fence rows and pastures, windbreaks, stone fences and stone piles, and rock ledges.

In the older-settled portions of the barberry eradication area, where cultivated barberries were introduced 50 to 100 years ago, many more barberry seeds have been scattered than in the newly-settled areas. In some of the communities in these older-settled States, escaped bushes and seedlings may be found on almost every farm. Areas of escaped bushes and seedlings increase in size as the scattered seedlings grow into bushes and produce seeds which are scattered in turn. In this way, seeds may be sown over a whole township or even a larger area in the course of years.

Areas of escaped barberries are brought under control as soon as practicable after they are located. All fruiting bushes and all bushes likely to fruit within three years usually are destroyed in the first treatment. If the area is small, all findable seedlings also are destroyed. If the area is large, destruction of small seedlings sometimes is delayed until the second treatment as a great many of the seedlings may succumb in the meantime to adverse weather conditions. In eradicating barberries from areas of escaped bushes, seasonal conditions that make for easier survey and eradication are used to advantage.

The interval elapsing between treatments is not long enough for any seedling to grow into a fruiting bush. This interval varies from three to five years, or possibly longer, depending upon local conditions.

ERADICATION

Salt is recommended for use in killing barberries. Kerosene is used if more convenient to procure, or if the salt may be disturbed or eaten by animals. However, the action of the kerosene is very slow. Barberry bushes close to valuable plants or trees are dug or pulled, as the application of chemicals may damage the other plants. Seedlings and small bushes sometimes are pulled, as it often is more economical to do so than to treat them with chemicals. That chemical eradication is effective is shown by the few sprouts that develop where chemicals have been applied properly.

During the calendar year 1926, 558.67 tons of salt and 1,962 gallons of kerosene were applied to 212,222 barberry bushes and sprouting bushes, and 1,663,829 seedlings, on 2,555 properties. (Tables 13 and 15.)

A total of 928,838 bushes, sprouting bushes, and seedlings was dug or pulled from 1,602 properties in the course of the three surveys. These figures show a rather large number dug or pulled, but about half of these were small bushes pulled from one area of escaped bushes. The total number eradicated by all methods during the year was 2,804,839 bushes, sprouting bushes, and seedlings.

SUMMARIZED RESULTS

Summary for 1926

During the calendar year, about 20 counties were covered in the original survey (Tables 1 and 2) and approximately 42 counties were surveyed for the second time. (Tables 5 and 6.) Resurvey was made where necessary to eliminate sprouts and seedlings. (Tables 9 and 10.) Original bushes numbering 204,530 were found on 2,555 properties and 723,530 bushes were destroyed on 3,057 properties. (Tables 1 and 2.) These totals include 52,286 bushes found on the 1,412 properties in second survey. (Tables 5 and 6.) In resurvey 16,149 sprouting bushes were found and 16,504 were eradicated. (Tables 9 and 10.) Seedlings numbering 2,062,639 were found in the course of three surveys.

Summary for Nine Years

The first survey of cities, towns, and villages was nearly completed in the earlier years of this survey. The 13 States include 978 counties. In the nine years of the campaign from April 1, 1918, to December 31, 1926, an area equivalent to approximately 363 counties has been covered in the original or first survey. (Tables 3 and 4.) A few counties will not be surveyed because of late settlement, arid conditions, or location in National Forests. Of about 920 counties necessary to survey, there remain about 57 yet to cover in the first survey.

A second survey has been made in 209,28 counties. (Tables 7 and 8.) The resurvey has been continued on properties covered by original and second surveys, to the extent necessary to eliminate sprouts and seedlings. (Tables 11 and 12.) A resurvey has been combined with the second survey in a large number of counties.

Original bushes numbering 6,705,423 have been located on 75,005 properties. Of these, 6,636,594 bushes have been destroyed on 74,686 properties. In resurvey 306,804 sprouting bushes were found on 13,451 properties and 306,238 sprouting bushes were destroyed from 13,438 properties. In all surveys 7,371,627 seedlings were found and 7,363,266 were destroyed. These numbers include 117,377 bushes and 163,661 seedlings found, and 117,856 bushes and 163,661 seedlings destroyed, on second survey. This makes a grand total of 14,383,354 bushes, sprouting bushes and seedlings found, and 14,361,093 bushes, sprouting bushes and seedlings destroyed, in all surveys of the entire campaign. These data for the nine years are summarized in tables 3, 4, 7, 8, 11, 12, 14, 16, and 17.

Credit is hereby gladly given to State Leaders, collaborators, and agents who have supplied data, and to Mrs. M. S. Koepfle, who has compiled, summarized, and tabulated them.

ORIGINAL SURVEY, PROPERTIES, January 1 to December 31, 1926

Table 1. Data showing, by States, the number of properties on which barberry bushes were found and destroyed in all surveys, and the number of properties upon which seedlings were found and destroyed in the original and second surveys during the calendar year January 1 to December 31, 1926

State	Number of : Number of properties on which			Total number of proper-			Number of properties on which		
	counties :	bushes were found	total in :	ties cleared of bushes	Found :	Destroyed	seedlings were -	Dug :	Treated: Total
	covered in:	In country	cities :	Dug :	Treated: Total	Found :	Dug :	Treated: Total	
	original :	In cities: Having :	and :	and :					
	survey :	and towns: escaped: Total :	country :	country :					
Colorado	0:	14:	10:	21:	35:	7:	0:	7:	7
Illinois	4.15:	333:	218:	367:	700:	413:	287:	700:	67 55: 12: 67 51
Indiana	0:	85:	22:	69:	154:	95:	60:	155:	20
Iowa	0:	53:	93:	162:	220:	64:	156:	220:	29
Michigan	2.10:	36:	232:	287:	323:	32:	746:	773:	125
Minnesota	0:	15:	43:	94:	109:	24:	85:	109:	20
Montana	6.83:	10:	16:	22:	32:	14:	14:	28:	3
Nebraska	0:	19:	26:	91:	110:	34:	92:	126:	11
North Dakota	0:	14:	0:	17:	31:	7:	24:	31:	2
Ohio	6.80:	94:	235:	364:	453:	205:	278:	433:	131
South Dakota	0:	14:	29:	74:	83:	12:	76:	83:	4
Wisconsin	0:	42:	220:	253:	295:	51:	223:	304:	13
Wyoming	0:	0:	0:	0:	0:	0:	0:	0:	0
Total	19.83:	734:	1,149:	1,821:	2,555:	987:	2,070:	3,057:	471 137: 340: 477

ORIGINAL SURVEY, BUSHES AND SEEDLINGS, JANUARY 1 to DECEMBER 31, 1926

Table 2. Data showing, by States, the number of barberry bushes found and destroyed in all surveys, and the number of seedlings found and destroyed in original and second surveys during the calendar year January 1 to December 31, 1926

State	Number of bushes found--			Number of bushes destroyed:			Number of seedlings			
	In cities: and towns:	In country: Escaped:	Total	Total	Dug	Treated	Total	Found	Destroyed	
									Dug	Treated
Colorado	84	251	287	371	9	363	372	606	0	606
Illinois	995	39,694	40,486	41,481	13,051	28,430	41,481	26,794	25,336	1,458
Indiana	351	262	440	791	1,231	408	1,639	1,461	1,291	120
Iowa	1,669	6,633	8,646	10,315	857	9,458	10,315	109,594	103	109,491
Michigan	159	73,565	85,345	85,504	275	96,096	96,371	1,323,226	32,235	1,290,991
Minnesota	216	2,746	3,294	3,510	242	3,268	3,510	3,021	136	2,885
Montana	72	97	293	365	75	281	356	223	183	35
Nebraska	34	1,557	2,161	2,195	494	1,765	2,259	1,066	1,026	41
North Dakota	69	0	330	399	215	184	399	27	0	27
Ohio	1,509	36,763	37,817	39,326	4,203	39,871	44,074	174,367	2,971	172,586
South Dakota	76	365	665	741	74	667	741	143	1	142
Wisconsin	245	18,998	19,287	19,532	497	24,775	522,063	17,209	5,843	12,366
Wyoming	0	0	0	0	0	0	0	0	0	0
Total	5,479	180,931	199,051	204,530	518,014	205,566	723,580	1,657,737	69,130	1,590,743
										1,659,878

ORIGINAL SURVEY, PROPERTIES, April 1, 1918, to December 31, 1926

Table 3. Data showing, by States, the number of properties on which barberry bushes were found and destroyed in all surveys, and the number of properties upon which seedlings were found and destroyed in original and second surveys, April 1, 1918, to December 31, 1926

State	Number of properties on which counties : bushes were found--		Total number of prop- ties cleared of bushes :		Number of properties on which seedlings were--							
	covered by: original survey	In cities: and towns: survey	Total in: cities and country	Dug Treated:	Found Dug	Destroyed Treated:						
Colorado	31.49:	1,560:	70:	186:	1,746:	1,653:	92:	1,745:	7:	0:	7:	7
Illinois	74.00:	10,956:	1,276:	3,291:	14,247:	12,559:	1,688:	14,247:	260:	202:	58:	260
Indiana	92.00:	3,698:	363:	1,322:	5,020:	4,499:	501:	5,000:	67:	32:	34:	66
Iowa	99.00:	7,108:	856:	2,857:	9,965:	9,095:	868:	9,963:	90:	37:	53:	90
Michigan	63.40:	5,121:	2,011:	5,718:	10,839:	9,017:	1,602:	10,619:	602:	426:	176:	602
Minesota	87.00:	3,178:	551:	2,117:	5,295:	4,940:	355:	5,295:	399:	361:	38:	399
Montana	43.16:	210:	35:	117:	327:	271:	49:	320:	13:	9:	4:	13
Nebraska	93.00:	3,213:	151:	832:	4,045:	3,693:	352:	4,045:	35:	18:	17:	35
North Dakota	53.00:	549:	1:	347:	896:	771:	125:	896:	4:	1:	3:	4
Ohio	78.65:	7,828:	1,101:	3,212:	11,040:	9,769:	1,247:	11,016:	324:	55:	269:	324
South Dakota	69.00:	504:	149:	696:	1,200:	819:	381:	1,200:	95:	85:	10:	95
Wisconsin	71.00:	7,044:	1,480:	3,249:	10,293:	9,059:	1,193:	10,252:	224:	100:	116:	216
Wyoming	8.12:	75:	1:	15:	90:	86:	2:	88:	0:	0:	0:	0
Total	\$62,82:	51,044:	\$,047:	23,961:	75,005:	66,231:	\$,455:	74,686:	2,120:	1,326:	785:	2,111

Table 4. Data showing, by States, the number of barberry bushes found and destroyed in all surveys, and the number of seedlings found and destroyed in original and second surveys, April 1, 1918, to December 31, 1926

State	Number of bushes found--			Number of bushes destroyed:			Number of seedlings--				
	In cities: and towns:	In country Escaped	Total	Dug	Treated:	Total	Found	Dug	Destroyed Treated	Total	
Colorado	19,681:	2,912:	5,124:	24,805:	23,917:	884:	24,801:	0:	606:	606	
Illinois	113,381:	213,162:	257,056:	370,437:	193,948:	176,489:	370,437:1,	46,229:1,	519,340:1,	565,569	
Indiana	77,821:	104,125:	119,365:	197,186:	99,045:	98,100:	197,145:	2,268:	7,759:	10,027	
Iowa	651,077:	65,976:	153,035:	804,162:	774,337:	29,519:	804,156:	1,838:	114,473:	116,311	
Michigan	54,143:	431,499:	513,061:	567,204:	367,123:	183,031:	555,154:2,	893,793:1,	366,277:2,	260,070	
Minnesota	592,708:	84,639:	196,758:	789,466:	780,820:	8,646:	789,466:	19,842:	5,483:	25,325	
Montana	6,985:	1,958:	4,673:	11,653:	10,097:	1,359:	11,456:	2,210:	1,775:	3,985	
Nebraska	73,119:	7,955:	24,592:	97,711:	91,165:	6,546:	97,711:	3,876:	6,370:	10,246	
North Dakota:	14,548:	150:	8,265:	22,813:	19,350:	2,963:	22,813:	150:	33:	183	
Ohio	219,624:	92,738:	111,844:	331,468:	250,367:	80,436:	330,803:	12,757:	329,861:	342,618	
South Dakota:	23,746:	21,243:	36,806:	60,552:	49,035:	11,467:	60,552:	16,865:	821:	17,686	
Wisconsin	281,243:3,	130,314:3,	142,542:3,	423,785:3,	350,377:	67,716:3,	418,093:	29,556:	30,207:	59,763	
Wyoming	3,947:	1:	229:	4,176:	3,972:	35:	4,007:	0:	0:	0	
Total	2,132,023:4,	156,722:4,	573,400:6,	705,423:6,	614,103:672,	491:6,	686,594:4,	415,725:1,	029,384:3,	383,005:4,	412,389

SECOND SURVEY, PROPERTIES, JANUARY 1 TO DECEMBER 31, 1926

Table 5. Data showing, by States, the number of properties on which barberry bushes and seedlings were found and destroyed on second survey in the barberry eradication campaign during the calendar year January 1 to December 31, 1926

State	Number of : counties: : surveyed:	Number of properties on which : bushes were found--		Total number of proper- : ties cleared of bushes		Number of properties on which : seedlings were--		Destroyed			
		In country	Total in : cities : and : country:	Dug	Treated	Total	Found	Dug	Treated	Total	
Colorado	1.05:	13:	7:	17:	30:	5:	25:	30:	7:	0:	7:
Illinois	4.21:	247:	211:	316:	563:	315:	248:	563:	60:	48:	12:
Indiana	2.20:	69:	8:	43:	112:	73:	36:	109:	14:	9:	5:
Iowa	7.63:	33:	56:	108:	141:	36:	105:	141:	14:	4:	10:
Michigan	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:
Minnesota	6.60:	9:	27:	64:	73:	16:	57:	73:	8:	4:	4:
Montana	1.40:	2:	4:	5:	7:	4:	2:	6:	1:	1:	0:
Nebraska	6.50:	19:	24:	83:	102:	29:	81:	110:	11:	9:	3:
North Dakota	3.60:	8:	0:	9:	17:	7:	10:	17:	2:	0:	2:
Ohio	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:
South Dakota	6.37:	14:	29:	74:	88:	12:	76:	88:	4:	1:	3:
Wisconsin	1.81:	42:	205:	237:	279:	73:	207:	280:	48:	13:	35:
Wyoming	.60:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:
Total	41.97:	456:	571:	956:	1,412:	570:	847:	1,417:	169:	89:	81:
											170

Table 6. Data showing, by States, the number of barberry bushes and seedlings found and destroyed on second survey in the barberry eradication campaign during the calendar year January 1 to December 31, 1926

[illegible]

SECOND SURVEY, PROPERTIES, January 1, 1922, to December 31, 1926

Table 7. Data showing, by States, the number of properties on which barberry bushes and seedlings were found and destroyed on second survey in barberry eradication campaign from January 1, 1922, to December 31, 1926

State	Number of properties on which bushes were found--			Total number of proper- ties cleared of bushes			Number of properties on which seedlings were--					
	Number of counties	In country	Total in cities & country	Dug	Treated	Total	Found	Dug	Treated	Total		
Colorado	21,09:	13:	23:	47:	60:	9:	51:	60:	7:	0:	7:	7
Illinois	4,85:	322:	242:	375:	697:	421:	276:	697:	85:	72:	13:	85
Indiana	9,35:	188:	13:	98:	286:	202:	76:	278:	18:	11:	7:	18
Iowa	21,63:	74:	178:	326:	400:	123:	276:	399:	40:	7:	33:	40
Michigan	3,00:	49:	38:	75:	124:	78:	46:	124:	4:	4:	0:	4
Minnesota	46,85:	74:	116:	383:	457:	182:	275:	457:	43:	15:	28:	43
Montana	10,13:	2:	4:	6:	8:	5:	2:	7:	1:	1:	0:	1
Nebraska	22,00:	67:	87:	263:	330:	82:	248:	330:	28:	17:	11:	28
North Dakota	29,65:	25:	0:	67:	92:	26:	66:	92:	3:	0:	3:	3
Ohio	2,00:	18:	0:	7:	25:	24:	1:	25:	0:	0:	0:	0
South Dakota	25,87:	43:	40:	167:	210:	33:	177:	210:	8:	5:	3:	8
Wisconsin	11,76:	201:	466:	615:	816:	340:	476:	816:	102:	52:	50:	102
Wyoming	1,10:	1:	0:	1:	2:	1:	1:	2:	0:	0:	0:	0
Total	209,28:	1,077:	1,207:	2,430:	3,507:	1,526:	1,971:	3,497:	339:	184:	155:	339

SECOND SURVEY, BUSHES AND SEEDLINGS, January 1, 1922, to December 31, 1926

Table 8. Data showing, by States, the number of barberry bushes and seedlings found and destroyed on second survey in the barberry eradication campaign from January 1, 1922, to December 31, 1926

State	Number of bushes found--			Number of bushes destroyed			Number of seedlings--		
	In cities:	In country	Total	Dug	Treated	Total	Found	Dug	Destroyed
Colorado	81:	80:	144:	225:	13:	212:	225:	606:	606:
Illinois	1,216:	84,645:	85,789:	87,005:	18,534:	68,471:	87,005:	18,800:	17,833:
Indiana	672:	55:	308:	980:	677:	285:	962:	1,327:	1,327:
Iowa	532:	2,277:	4,632:	5,164:	1,073:	4,089:	5,162:	105,522:	104,455:
Michigan	96:	339:	438:	534:	262:	272:	534:	240:	240:
Minnesota	458:	2,006:	4,823:	5,281:	1,758:	3,523:	5,281:	3,627:	536:
Montana	2:	77:	82:	84:	74:	9:	83:	29:	29:
Nebraska	506:	2,609:	5,032:	5,538:	1,235:	4,303:	5,538:	4,496:	3,476:
North Dakota	145:	0:	1,703:	1,848:	315:	1,533:	1,848:	30:	0:
Ohio	59:	0:	31:	90:	88:	2:	90:	0:	0:
South Dakota	344:	334:	1,759:	2,103:	324:	1,779:	2,103:	828:	686:
Wisconsin	741:	7,535:	8,250:	8,991:	3,218:	5,773:	8,991:	28,156:	19,059:
Wyoming	1:	0:	33:	34:	1:	33:	34:	0:	0:
Total	4,853:	99,957:	113,024:	117,877:	27,572:	90,284:	117,856:	163,661:	43,381:
								120,280:	163,661

Table 9. Data showing, by States, the number of properties on which sprouting bushes and seedlings were found and destroyed on resurvey in the barberry eradication campaign during the calendar year January 1 to December 31, 1926

State	Number of properties on which sprouting bushes were found--			Total number of properties : cleared of sprouting bushes:			Number of properties on which seedlings were--			Total	
	In cities: and towns:	In country bushes :	Total in: cities & escaped: bushes :	Dug	Treated	Total	Destroyed				
							Dug	Treated			
Colorado	11:	9:	16:	27:	4:	24:	28:	10:	1:	9:	10
Illinois	38:	86:	115:	153:	81:	72:	153:	91:	80:	11:	91
Indiana	16:	21:	46:	62:	38:	23:	61:	9:	3:	5:	8
Iowa	25:	43:	123:	148:	49:	99:	148:	63:	27:	36:	63
Michigan	10:	2:	2:	12:	8:	4:	12:	2:	2:	0:	2
Minnesota	29:	72:	137:	166:	55:	111:	166:	50:	30:	20:	50
Montana	6:	3:	4:	10:	7:	3:	10:	9:	8:	1:	9
Nebraska	14:	7:	48:	62:	25:	39:	64:	3:	2:	1:	3
North Dakota	48:	0:	44:	92:	20:	72:	92:	5:	0:	5:	5
Ohio	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0
South Dakota	1:	2:	7:	8:	1:	7:	8:	5:	4:	1:	5
Wisconsin	19:	17:	34:	53:	22:	31:	53:	22:	11:	11:	22
Wyoming	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0
Total	217:	262:	576:	793:	310:	485:	795:	269:	168:	100:	268

Table 10. Data showing, by States, the number of sprouting bushes and seedlings found and destroyed on resurvey in the barberry eradication campaign during the calendar year January 1 to December 31, 1926

State	Number of sprouting bushes found--			Number of sprouting bushes destroyed			Number of seedlings--				
	In cities:	In country	Total	Dug	Treated	Total	Found	Dug	Destroyed		
Colorado	31:	43:	74:	105:	7:	99:	106:	216:	4:	212:	216
Illinois	157:	3,036:	6,134:	6,291:	3,466:	2,825:	6,291:	372,077:	321,800:	50,277:	372,077
Indiana	34:	163:	220:	254:	93:	160:	253:	806:	431:	350:	781
Iowa	233:	400:	6,082:	6,315:	5,590:	1,077:	6,667:	14,801:	1,240:	13,561:	14,801
Michigan	13:	25:	25:	38:	25:	13:	38:	5:	5:	0:	5
Minnesota	111:	969:	1,372:	1,483:	331:	1,152:	1,483:	2,065:	1,379:	686:	2,065
Montana	15:	2:	18:	33:	19:	14:	33:	60:	37:	23:	60
Nebraska	162:	124:	579:	741:	160:	584:	744:	367:	267:	100:	367
North Dakota	220:	0:	210:	430:	85:	345:	430:	99:	0:	99:	99
Ohio	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0
South Dakota	1:	2:	35:	36:	1:	35:	36:	40:	32:	8:	40
Wisconsin	80:	293:	343:	423:	71:	352:	423:	14,416:	6,651:	7,765:	14,416
Wyoming	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0
Total	1,057:	5,057:	15,092:	16,149:	9,848:	6,656:	16,504:	404,952:	331,846:	73,081:	404,927

RESURVEY, PROPERTIES, April 1, 1918, to December 31, 1926

Table 11. Data showing, by States, the number of properties on which sprouting bushes and seedlings were found and destroyed on resurvey in the barberry eradication campaign from April 1, 1918, to December 31, 1926

State	Number of properties on which sprout--: Total number of properties: Number of properties on which				ing bushes were found--: cleared of sprouting bushes: seedlings were--				Destroyed			
	In cities:	In country	Total in	Dug	Treated	Total	Found	Dug	Treated	Total		
	and towns:	escaped:	and									
	bushes:		country									
Colorado	1,442:	114:	193:	1,635:	1,416:	219:	1,635:	102:	19:	83:	102	
Illinois	442:	395:	767:	1,209:	572:	637:	1,209:	387:	316:	71:	387	
Indiana	178:	136:	266:	444:	305:	134:	439:	48:	14:	33:	47	
Iowa	343:	329:	1,011:	1,354:	703:	646:	1,354:	275:	142:	133:	275	
Michigan	146:	114:	289:	435:	375:	60:	435:	191:	187:	4:	191	
Minnesota	736:	612:	1,295:	2,031:	1,630:	401:	2,031:	2,241:	2,114:	127:	2,241	
Montana	123:	6:	56:	179:	163:	16:	179:	23:	20:	3:	23	
Nebraska	211:	26:	413:	624:	363:	261:	624:	33:	14:	19:	33	
North Dakota	295:	0:	223:	518:	243:	275:	518:	6:	0:	6:	6	
Ohio	1,421:	281:	1,021:	2,442:	2,104:	338:	2,442:	716:	551:	165:	716	
South Dakota	341:	35:	338:	679:	513:	166:	679:	79:	36:	43:	79	
Wisconsin	913:	655:	946:	1,859:	1,344:	511:	1,855:	274:	168:	106:	274	
Wyoming	32:	0:	10:	42:	31:	7:	33:	7:	7:	0:	7	
Total	6,623:	2,703:	6,823:	13,451:	9,767:	3,671:	13,438:	4,382:	3,588:	793:	4,381	

RESURVEY, SPROUTING BUSHES AND SEEDLINGS, April 1, 1918, to December 31, 1926

Table 12. Data showing, by States, the number of sprouting bushes and seedlings found and destroyed on resurvey in the barberry eradication campaign from April 1, 1918, to December 31, 1926

State	Number of sprouting bushes found--			Number of sprouting bushes:			Number of seedlings--		
	In cities:	In country	Total	Destroyed--	Dug	Treated	Found	Dug	Treated
	and towns:	Escaped	Total	Total	Dug	Treated	Total	Dug	Treated
Colorado	3,827:	2,023:	3,160:	6,987:	5,148:	1,839:	6,987:	3,672:	712:
Illinois	4,816:	7,902:	16,626:	21,442:	9,923:	11,519:	574,840:	399,043:	175,797:
Indiana	1,529:	16,871:	18,310:	19,839:	17,592:	1,941:	4,296:	719:	3,552:
Iowa	4,009:	8,205:	23,942:	27,951:	15,780:	12,171:	57,648:	28,774:	28,874:
Michigan	524:	1,194:	2,418:	2,942:	2,231:	711:	2,942:	547,349:	60,085:
Minnesota	14,039:	17,413:	36,832:	50,871:	40,690:	10,181:	50,871:	28,097:	23,695:
Montana	3,557:	5:	1,647:	5,204:	5,070:	134:	5,204:	1,069:	670:
Nebraska	6,172:	249:	10,415:	16,587:	12,564:	4,023:	16,587:	1,243:	1,817:
North Dakota	854:	0:	1,260:	2,114:	281:	1,833:	2,114:	0:	100:
Ohio	5,666:	8,046:	12,276:	17,942:	13,071:	4,871:	362,585:	111,527:	251,058:
South Dakota	20,980:	5,284:	22,045:	43,025:	36,618:	6,407:	43,025:	6,494:	2,611:
Wisconsin	11,244:	75,129:	80,081:	91,325:	19,440:	71,704:	91,144:1,303,943:	136,941:1,167,002:1,303,943	0:
Wyoming	546:	0:	29:	575:	475:	21:	496:	53:	0:
Total	77,763:	142,321:	229,041:	306,804:	178,883:	127,355:	306,238:2,955,902:1,237,656:1,718,221:2,955,877		

CHEMICAL TREATMENT, 1926

Table 13. Data showing, by States, the number of properties on which barberry bushes and sprouting barberry bushes were treated with chemicals, and the number of bushes, sprouting bushes, and seedlings treated from January 1 to December 31, 1926

State	Number treated--												Total
	With Salt		With sodium arsenite		With Kerosene								
	Proper- ties	Bushes	Proper- ties	Seed- lings	Proper- ties	Seed- lings	Proper- ties	Bushes	Seed- lings	Proper- ties	Bushes		
Colorado	48:	428:	818:	0:	0:	5:	34:	0:	53:	462:	818		
Illinois	359:	31,255:	51,735:	0:	0:	0:	0:	0:	359:	31,255:	51,735		
Indiana	66:	462:	427:	0:	0:	17:	106:	43:	83:	568:	470		
Iowa	247:	9,521:	123,052:	0:	0:	8:	1,014:	0:	255:	10,535:	123,052		
Michigan	750:	96,109:	1,290,991:	0:	0:	0:	0:	0:	750:	96,109:	1,290,991		
Minnesota	194:	4,401:	3,571:	0:	0:	2:	19:	0:	196:	4,420:	3,571		
Montana	17:	295:	58:	0:	0:	0:	0:	0:	17:	295:	58		
Nebraska	46:	1,281:	40:	0:	0:	85:	1,168:	101:	131:	2,349:	141		
North Dakota	96:	529:	126:	0:	0:	0:	0:	0:	96:	529:	126		
Ohio	265:	39,799:	168,162:	0:	0:	13:	72:	4,424:	278:	39,871:	172,586		
South Dakota	83:	702:	150:	0:	0:	0:	0:	0:	83:	702:	150		
Wisconsin	254:	25,127:	20,131:	0:	0:	0:	0:	0:	254:	25,127:	20,131		
Wyoming	0:	0:	0:	0:	0:	0:	0:	0:	0:	0:	0		
Total	2,425:	209,909:	1,659,261:	0:	0:	130:	2,313:	4,568:	2,555:	212,222:	1,663,829		

CHEMICAL TREATMENT, Sept. 1, 1921, to December 31, 1926

Table 14. Data showing, by States, the number of properties on which barberry bushes and sprouting barberry bushes were treated with chemicals, and the number of bushes, sprouting bushes, and seedlings treated from Sept. 1, 1921, to December 31, 1926

State	With salt			With sodium arsenite			With kerosene			Total		
	Proper- ties	Bushes	Seedlings	Proper- ties	Bushes	Seedlings	Proper- ties	Bushes	Seedlings	Proper- ties	Bushes	Seedlings
Colorado	303	2,640	3,566	0	0	0	8	83	0	311	2,723	3,566
Illinois	2,286	186,000	1,695,137	34	839	0	5	1,169	0	2,325	188,008	1,695,137
Indiana	618	99,935	11,268	0	0	0	17	106	43	635	100,041	11,311
Iowa	1,487	40,696	143,344	0	0	0	27	1,294	3	1,514	41,990	143,347
Michigan	1,286	117,961	1,298,813	239	8,594	29,911	137	62,187	97	1,662	188,742	1,426,362
Minnesota	727	18,698	29,046	25	85	102	4	44	30	756	18,827	29,178
Montana	65	1,493	2,445	0	0	0	0	0	0	65	1,493	2,445
Nebraska	435	8,219	7,483	0	0	0	178	2,350	704	613	10,569	8,187
North Dakota	379	4,729	133	21	67	0	0	0	0	400	4,796	133
Ohio	1,303	73,150	435,035	10	1,069	59,300	272	11,088	86	1,585	35,307	580,919
South Dakota	539	17,861	3,416	0	0	0	8	13	16	547	17,874	3,432
Wisconsin	1,353	133,595	1,195,507	350	5,824	1,702	1	1	0	1,704	139,420	1,197,209
Wyoming	9	56	0	0	0	0	0	0	0	9	56	0
Total	10,790	705,033	4,825,243	679	16,478	91,015	657	78,335	184,963	12,126	799,846	5,101,226

CHEMICALS, QUANTITIES USED, January 1, to December 31, 1926

Table 15. Data showing, by States, quantities of chemicals used in the barberry eradication campaign from January 1 to December 31, 1926

State	Salt (Tons)			Sodium arsenite (Gals.)			Kerosene (Gallons)		
	Furnished by--	Property:	State:	Conference:	Total	Conference:	Furnished by--	Property:	State:
	Owner	Agency:	P. G. Rust	U. S. D. A.		P. G. Rust	U. S. D. A.	Total	Owner
Colorado	0:	0:	0:	1.19:	1.19:	0:	0:	0:	49.0:
Illinois	0:	.91:	0:	46.80:	47.71:	0:	0:	0:	0
Indiana	.01:	0:	0:	2.40:	2.41:	0:	0:	0:	132.0
Iowa	.04:	0:	.54:	34.74:	35.32:	0:	0:	0:	314.5:
Michigan	0:	0:	0:	171.19:	171.19:	0:	0:	0:	0
Minnesota	.75:	0:	0:	11.92:	12.67:	0:	0:	0:	17.0
Montana	0:	0:	0:	.82:	.82:	0:	0:	0:	0
Nebraska	0:	0:	0:	2.94:	2.94:	0:	0:	0:	1.5:
North Dakota	2.50:	0:	0:	.20:	2.70:	0:	0:	0:	0
Ohio	.12:	207.07:	0:	14.32:	222.01:	0:	0:	0:	29.0:
South Dakota	.26:	0:	0:	4.99:	5.25:	0:	0:	0:	253.0
Wisconsin	0:	54.37:	0:	.09:	54.46:	0:	0:	0:	0
Wyoming	0:	0:	0:	0:	0:	0:	0:	0:	0
Total	3,68:	262.35:	.54:	292.10:	558.67:	0:	0:	0:	34.0:
									1,928.0:
									1,962.0

CHEMICALS, QUANTITIES USED, Sept: 1, 1921, to December 31, 1926

Table 16. Data showing, by States, quantities of chemicals used in the barberry eradication campaign from Sept: 1, 1921, to December 31, 1926

State	Salt (Tons)			Sodium Arsenite (Gallons)			Kerosene (Gallons)		
	Furnished by--			Furnished by--			Furnished by--		
	Property:Owner	State:agency	Conference:P. G. Rust	Conference:U.S.D.A.	Total:P. G. Rust	Total:U.S.D.A.	Owner:U.S.D.A.	Total	Total
Colorado	0:	0:	0:	5.16:	5.16:	0:	0:	80.0	80.0
Illinois	.75:	55.78:	31.00:	314.07:	401.60:	0:	0:	124.0	124.0
Indiana	.83:	0:	0:	58.45:	59.28:	0:	0:	132.0	132.0
Iowa	44.09:	0:	20.69:	92.58:	157.36:	0:	0:	404.25:	718.75
Michigan	.03:	0:	8.49:	257.33:	265.85:	175.6:	129.3:	304.9:	0:11,341.0
Minnesota	2.88:	.84:	9.21:	40.89:	53.82:	0:	23.25:	23.25:	27.0 ^a
Montana	.12:	0:	0:	5.05:	5.17:	0:	0:	0:	0
Nebraska	.13:	0:	8.55:	19.23:	27.91:	0:	0:	151.5:	2,597.5
North Dakota	14.63:	5.00:	0:	4.88:	24.51:	0:	7.0:	0:	2,749.0
Ohio	3.02:	342.48:	0:	23.37:	368.87:	16.2:	30.1:	46.3:	5,182.0*
South Dakota	14.37:	0:	17.85:	9.79:	42.01:	0:	0:	0:	22.0
Wisconsin	.25:	174.82:	70.00:	25.29:	270.36:	408.0:	190.0:	598.0:	3,375 ^b
Wyoming	.05:	0:	0:	.16:	.21:	0:	0:	0:	0
Total	81.15:	578.92:	165.79:	856.25:	1,682.11:	599.8:	456.65:	1,056.45:	5,737.75:
									16,276.375
									22,014.125

^a 10 gallons of drip oil

^b carbon bisulphide

* 4,903 gallons furnished by State

GRAND SUMMARY, ORIGINAL BUSHES, SPROUTING BUSHES, AND SEEDLINGS, 1918 to 1926

Table 17. Data showing, by States, the number of bushes, sprouting bushes, and seedlings found and destroyed in all surveys in the barberry eradication campaign, April 1, 1918, to December 31, 1926

State	Original bushes		Sprouting bushes		Seedlings		Grand Total	
	Found	Destroyed	Found	Destroyed	Found	Destroyed	Found	Destroyed
Colorado	24,805	24,801	6,987	6,987	4,278	4,278	36,070	36,066
Illinois	370,437	370,437	21,442	21,442	2,140,409	2,140,409	2,532,288	2,532,288
Indiana	197,186	197,145	19,839	19,533	14,373	14,298	231,398	230,976
Iowa	804,162	804,156	27,951	27,951	173,959	173,959	1,006,072	1,006,066
Michigan	567,204	555,154	2,942	2,942	2,867,504	2,867,504	3,437,650	3,425,600
Minnesota	789,466	789,466	50,871	50,871	53,422	53,422	893,759	893,759
Montana	11,658	11,456	5,204	5,204	5,054	5,054	21,916	21,714
Nebraska	97,711	97,711	16,587	16,587	13,306	13,306	127,604	127,604
North Dakota	22,813	22,813	2,114	2,114	283	283	25,210	25,210
Ohio	331,468	330,803	17,942	17,942	705,203	705,203	1,054,613	1,053,948
South Dakota	60,552	60,552	43,025	43,025	26,791	26,791	130,368	130,368
Wisconsin	3,423,785	3,418,093	91,325	91,144	1,366,992	1,363,706	4,882,102	4,872,943
Wyoming	4,176	4,007	575	496	53	53	4,804	4,556
Total	6,705,423	6,686,594	306,804	306,238	7,371,627	7,368,266	14,383,854	14,361,098

